

# Le monoïde de 0-Rook, et autres monoïdes $\mathcal{T}$ -triviaux

Joël Gay

joint work with Florent Hivert

Université Paris-Sud, LRI & École Polytechnique, LIX

Conference for Ph. D students - April 28, 2017

# Contents

## 1 Green Relations

## 2 The 0-Hecke monoid

## 3 Representation theory of $\mathcal{J}$ -trivial monoid

## 4 The 0-rook monoid

## 5 Renner monoids

# Some basic definitions

## Group

## Monoid

## Semigroup

# Some basic definitions

## Group

## Monoid

## Semigroup

$(G, \cdot)$ ,  $\cdot$  intern and

- Associative
- Has a unit
- Each element has an inverse

# Some basic definitions

## Group

$(G, \cdot)$ ,  $\cdot$  intern and

- Associative
- Has a unit
- Each element has an inverse

## Monoid

$(M, \cdot)$ ,  $\cdot$  intern and

- Associative
- Has a unit

## Semigroup

# Some basic definitions

## Group

$(G, \cdot)$ ,  $\cdot$  intern and

- Associative
- Has a unit
- Each element has an inverse

## Monoid

$(M, \cdot)$ ,  $\cdot$  intern and

- Associative
- Has a unit

## Semigroup

$(S, \cdot)$ ,  $\cdot$  intern and

- Associative

# Some basic definitions

## Group

$(G, \cdot), \cdot$  intern and

- Associative
- Has a unit
- Each element has an inverse

**Representation** :  $(\rho, V)$  with

- $V$   $\mathbb{C}$ -vectorial space
- $\rho : G \rightarrow GL(V)$  group morphism
- if  $\dim V < \infty$  :  $GL(V) \cong GL_n(\mathbb{C})$

## Monoid

$(M, \cdot), \cdot$  intern and

- Associative
- Has a unit

## Semigroup

$(S, \cdot), \cdot$  intern and

- Associative

# Some basic definitions

## Group

$(G, \cdot)$ ,  $\cdot$  intern and

- Associative
- Has a unit
- Each element has an inverse

**Representation** :  $(\rho, V)$  with

- $V$   $\mathbb{C}$ -vectorial space
- $\rho : G \rightarrow GL(V)$  group morphism
- if  $\dim V < \infty$  :  $GL(V) \cong GL_n(\mathbb{C})$

## Monoid

$(M, \cdot)$ ,  $\cdot$  intern and

- Associative
- Has a unit

**Representation** :  $(\rho, V)$  with

- $V$   $\mathbb{C}$ -vectorial space
- $\rho : G \rightarrow \text{End}(V)$  monoid morphism
- if  $\dim V < \infty$  :  $\text{End}(V) \cong M_n(\mathbb{C})$

## Semigroup

$(S, \cdot)$ ,  $\cdot$  intern and

- Associative



# Some basic definitions

## Group

$(G, \cdot)$ ,  $\cdot$  intern and

- Associative
- Has a unit
- Each element has an inverse

**Representation** :  $(\rho, V)$  with

- $V$   $\mathbb{C}$ -vectorial space
- $\rho : G \rightarrow GL(V)$  group morphism
- if  $\dim V < \infty$  :  $GL(V) \cong GL_n(\mathbb{C})$

## Monoid

$(M, \cdot)$ ,  $\cdot$  intern and

- Associative
- Has a unit

**Representation** :  $(\rho, V)$  with

- $V$   $\mathbb{C}$ -vectorial space
- $\rho : G \rightarrow \text{End}(V)$  monoid morphism
- if  $\dim V < \infty$  :  $\text{End}(V) \cong M_n(\mathbb{C})$

## Semigroup

$(S, \cdot)$ ,  $\cdot$  intern and

- Associative

## Group algebra $\mathbb{C}G$

# Some basic definitions

## Group

$(G, \cdot)$ ,  $\cdot$  intern and

- Associative
- Has a unit
- Each element has an inverse

**Representation** :  $(\rho, V)$  with

- $V$   $\mathbb{C}$ -vectorial space
- $\rho : G \rightarrow GL(V)$  group morphism
- if  $\dim V < \infty$  :  $GL(V) \cong GL_n(\mathbb{C})$

**Group algebra**  $\mathbb{C}G$

## Monoid

$(M, \cdot)$ ,  $\cdot$  intern and

- Associative
- Has a unit

**Representation** :  $(\rho, V)$  with

- $V$   $\mathbb{C}$ -vectorial space
- $\rho : G \rightarrow \text{End}(V)$  monoid morphism
- if  $\dim V < \infty$  :  $\text{End}(V) \cong M_n(\mathbb{C})$

**Monoid algebra**  $\mathbb{C}M$

# Preorders and equivalence relations

## Definition (Green)

Let  $M$  a monoid,  $x, y \in M$ . We say :

- $x \leq_{\mathcal{R}} y$  iff  $xM \subseteq yM$
- $x \leq_{\mathcal{L}} y$  iff  $Mx \subseteq My$
- $x \leq_{\mathcal{J}} y$  iff  $MxM \subseteq MyM$

# Preorders and equivalence relations

## Definition (Green)

Let  $M$  a monoid,  $x, y \in M$ . We say :

- $x \leq_{\mathcal{R}} y$  iff  $xM \subseteq yM$  iff  $x = yu$  for some  $u \in M$ .
- $x \leq_{\mathcal{L}} y$  iff  $Mx \subseteq My$  iff  $x = uy$  for some  $u \in M$ .
- $x \leq_{\mathcal{J}} y$  iff  $MxM \subseteq MyM$  iff  $x = uyv$  for some  $u, v \in M$ .

# Preorders and equivalence relations

## Definition (Green)

Let  $M$  a monoid,  $x, y \in M$ . We say :

- $x \leq_{\mathcal{R}} y$  iff  $xM \subseteq yM$  iff  $x = yu$  for some  $u \in M$ .
- $x \leq_{\mathcal{L}} y$  iff  $Mx \subseteq My$  iff  $x = uy$  for some  $u \in M$ .
- $x \leq_{\mathcal{J}} y$  iff  $MxM \subseteq MyM$  iff  $x = uyv$  for some  $u, v \in M$ .

Warning : 1 on top!

# Preorders and equivalence relations

## Definition (Green)

Let  $M$  a monoid,  $x, y \in M$ . We say :

- $x\mathcal{R}y$  iff  $xM = yM$ .
- $x\mathcal{L}y$  iff  $Mx = My$ .
- $x\mathcal{J}y$  iff  $MxM = MyM$ .

# $\mathcal{K}$ -triviality

## Definition

A monoid is  $\mathcal{K}$ -trivial ( $\mathcal{K} \in \{\mathcal{R}, \mathcal{L}, \mathcal{J}\}$ ) if all its  $\mathcal{K}$ -classes are of cardinality one.

# Ordered monoids

## Definition (Green)

Let  $M$  a monoid with  $\leq$  a partial order.  $M$  is called :

- right ordered iff  $xy \leq x$  for all  $x, y \in M$ .
- left ordered iff  $yx \leq x$  for all  $x, y \in M$ .
- left-right ordered iff  $xy \leq x$  and  $yx \leq x$  for all  $x, y \in M$ .



# Ordered monoids

## Proposition (Green)

Let  $M$  a monoid. It is :

- right ordered iff  $\mathcal{R}$ -trivial.
- left ordered iff  $\mathcal{L}$ -trivial.
- left-right ordered iff  $\mathcal{J}$ -trivial

# Ordered monoids

## Proposition (Green)

Let  $M$  a monoid. It is :


- right ordered iff  $\mathcal{R}$ -trivial.
- left ordered iff  $\mathcal{L}$ -trivial.
- left-right ordered iff  $\mathcal{J}$ -trivial

$\mathcal{K}$ -triviality can be seen in Cayley graph

# Example : Cayley-graph and $\mathcal{K}$ -triviality

$$M = \langle a, b \rangle$$


$$a^2 = a ; b^2 = b ; aba = ab ; bab = ba$$



# Example : Cayley-graph and $\mathcal{K}$ -triviality

$$M = \langle a, b \rangle$$


$$a^2 = a ; b^2 = b ; aba = ab ; bab = ba$$



# Example : Cayley-graph and $\mathcal{K}$ -triviality


$$M = \langle a, b \rangle$$

$$a^2 = a ; b^2 = b ; aba = ab ; bab = ba$$




# Aperiodic monoid

Let  $M$  a monoid,  $x \in M$ .



# Aperiodic monoid

Let  $M$  a monoid,  $x \in M$ .




## Definition

$M$  is said *aperiodic* is  $\forall x \in M, \exists n, x^n = x^{n+1}$ .


Notation  $x^\omega := x^n$  idempotent

$E(M)$  : set of idempotents of  $M$ .



# Aperiodic monoid

Let  $M$  a monoid,  $x \in M$ .




## Definition

$M$  is said *aperiodic* is  $\forall x \in M, \exists n, x^n = x^{n+1}$ .

Notation  $x^\omega := x^n$  idempotent

$E(M)$  : set of idempotents of  $M$ .



$\mathcal{K}$ -trivial  $\Rightarrow$  aperiodic.



# Idempotents and representation of finite semigroup

Let  $S$  a finite semigroup.

- $e \in S$  idempotent  $\Rightarrow eSe$  monoid with identity  $e$ .  
 $G_e$  group of units.

# Idempotents and representation of finite semigroup

Let  $S$  a finite semigroup.

- $e \in S$  idempotent  $\Rightarrow eSe$  monoid with identity  $e$ .  
 $G_e$  group of units.
- If  $e, f$  idempotents.  $e\mathcal{J}f \Rightarrow G_e \cong G_f$ .

# Idempotents and representation of finite semigroup

Let  $S$  a finite semigroup.

- $e \in S$  idempotent  $\Rightarrow eSe$  monoid with identity  $e$ .  
 $G_e$  group of units.
- If  $e, f$  idempotents.  $e\mathcal{J}f \Rightarrow G_e \cong G_f$ .
- A  $\mathcal{J}$ -class  $J$  is said *regular* if it contains an idempotent.  
If  $e \in J$  then  $G_e = eSe \cap J$ .

# Idempotents and representation of finite semigroup

## Theorem (Clifford, Munn, Ponizovskii)

*Let  $S$  a semigroup,  $K$  a field,  $E = \{e_J\}$  an idempotent transversal over the regular  $\mathcal{J}$ -classes  $J$  of  $S$ . Let  $G_J := G_{e_J}$ .*

*Then there is a bijection between irreducible representations of  $S$  and irreducible representations of the  $G_J$ .*

# Idempotents and representation of finite semigroup

## Theorem (Clifford, Munn, Ponizovskii)

*Let  $S$  a semigroup,  $K$  a field,  $E = \{e_J\}$  an idempotent transversal over the regular  $\mathcal{J}$ -classes  $J$  of  $S$ . Let  $G_J := G_{e_J}$ .*

*Then there is a bijection between irreducible representations of  $S$  and irreducible representations of the  $G_J$ .*

Remark : the aperiodic monoids is the class of monoids for which all  $G_J$  are trivial.

# Contents

1 Green Relations

2 The 0-Hecke monoid

3 Representation theory of  $\mathcal{J}$ -trivial monoid

4 The 0-rook monoid

5 Renner monoids

# Definition by presentation

$$\mathfrak{S}_n$$

Symmetric Group

# Definition by presentation

$$\mathfrak{S}_n \xleftarrow{q=1} \mathcal{H}_n(q)$$

Symmetric Group

Iwahori-Hecke  
algebra



# Definition by presentation

$$\mathfrak{S}_n \xleftarrow{q=1} \mathcal{H}_n(q) \xrightarrow{q=0} H_n^0$$

Symmetric Group

Iwahori-Hecke  
algebraHecke monoid at  
 $q = 0$

# Definition by presentation

$$\mathfrak{S}_n \xleftarrow{q=1} \mathcal{H}_n(q) \xrightarrow{q=0} H_n^0$$

Symmetric Group

Iwahori-Hecke  
algebraHecke monoid at  
 $q = 0$ 

$$s_i^2 = 1$$

$$T_i^2 = q1 + (q - 1)T_i$$

$$\pi_i^2 = \pi_i$$

$$s_{i+1}s_i s_{i+1} = s_i s_{i+1} s_i$$

$$T_{i+1}T_i T_{i+1} = T_i T_{i+1} T_i$$

$$\pi_{i+1}\pi_i \pi_{i+1} = \pi_i \pi_{i+1} \pi_i$$

$$s_i s_j = s_j s_i$$

$$T_i T_j = T_j T_i$$

$$\pi_i \pi_j = \pi_j \pi_i$$

# Definition by presentation

$$\mathfrak{S}_n \xleftarrow{q=1} \mathcal{H}_n(q) \xrightarrow{q=0} H_n^0$$

Symmetric Group

Iwahori-Hecke  
algebraHecke monoid at  
 $q = 0$ 

$$s_i^2 = 1$$

$$T_i^2 = q1 + (q - 1)T_i$$

$$\pi_i^2 = \pi_i$$

$$s_{i+1}s_i s_{i+1} = s_i s_{i+1} s_i$$

$$T_{i+1}T_i T_{i+1} = T_i T_{i+1} T_i$$

$$\pi_{i+1}\pi_i \pi_{i+1} = \pi_i \pi_{i+1} \pi_i$$

$$s_i s_j = s_j s_i$$

$$T_i T_j = T_j T_i$$

$$\pi_i \pi_j = \pi_j \pi_i$$

# Definition by presentation

$$\mathfrak{S}_n \xleftarrow{q=1} \mathcal{H}_n(q) \xrightarrow{q=0} H_n^0$$

Symmetric Group

Iwahori-Hecke  
algebraHecke monoid at  
 $q = 0$ 

$$s_i^2 = 1$$

$$T_i^2 = q1 + (q-1)T_i$$

$$\pi_i^2 = \pi_i$$

$$s_{i+1}s_i s_{i+1} = s_i s_{i+1} s_i \quad T_{i+1}T_i T_{i+1} = T_i T_{i+1} T_i$$

$$\pi_{i+1}\pi_i \pi_{i+1} = \pi_i \pi_{i+1} \pi_i$$

$$s_i s_j = s_j s_i$$

$$T_i T_j = T_j T_i$$

$$\pi_i \pi_j = \pi_j \pi_i$$

$$\begin{array}{c} \frown \\ s_i = T_i \end{array}$$

$$\begin{array}{c} \frown \\ \pi_i = T_i + 1 \end{array}$$

# Definition by action

Action on  $\mathfrak{S}_n$

Bubble sort algorithm :

$$(s_1 \dots s_n) \cdot \pi_i = \begin{cases} s_1 \dots s_{i-1} s_{i+1} s_i s_{i+2} \dots s_n & \text{if } s_i < s_{i+1}, \\ s_1 \dots s_n & \text{otherwise.} \end{cases}$$

# Definition by action

Action on  $\mathfrak{S}_n$

Bubble sort algorithm :

$$(s_1 \dots s_n) \cdot \pi_i = \begin{cases} s_1 \dots s_{i-1} s_{i+1} s_i s_{i+2} \dots s_n & \text{if } s_i < s_{i+1}, \\ s_1 \dots s_n & \text{otherwise.} \end{cases}$$

$$3726\textcolor{red}{1}45 \cdot \pi_5 = 3726\textcolor{red}{4}15$$

$$3726\textcolor{red}{4}15 \cdot \pi_5 = 3726\textcolor{red}{4}15$$

# Theorems for $H_n^0$

## Theorem

*The two definitions are equivalent.*

# Theorems for $H_n^0$

## Theorem

*The two definitions are equivalent.*

## Theorem

$$|H_n^0| = |\mathfrak{S}_n| = n!$$



# Theorems for $H_n^0$

## Theorem


*The two definitions are equivalent.*

## Theorem

$$|H_n^0| = |\mathfrak{S}_n| = n!$$

## Theorem

*An element of  $H_n^0$  is characterized by its action on the identity.*



# Theorems for $H_n^0$

## Theorem


*The two definitions are equivalent.*

## Theorem

$$|H_n^0| = |\mathfrak{S}_n| = n!$$

## Theorem

*An element of  $H_n^0$  is characterized by its action on the identity.*



# Theorems for $H_n^0$

## Theorem

*The two definitions are equivalent.*

## Theorem


$$|H_n^0| = |\mathfrak{S}_n| = n!$$

## Theorem


*An element of  $H_n^0$  is characterized by its action on the identity.*

## Theorem


*$H_n^0$  is  $\mathcal{T}$ -trivial*




# $\mathcal{T}$ -triviality : $H_n^0$ right Cayley graph



# $\mathcal{T}$ -triviality : $H_n^0$ left Cayley graph



# $\mathcal{J}$ -triviality : $H_n^0$ bisided Cayley graph



# Contents

- 1 Green Relations
- 2 The 0-Hecke monoid
- 3 Representation theory of  $\mathcal{J}$ -trivial monoid
- 4 The 0-rook monoid
- 5 Renner monoids

# Simple modules

$M$  is  $\mathcal{J}$ -trivial and  $x \in M$ .

## Definition

Let  $S_x = \langle \epsilon_x \rangle$  vector space, right action of  $y \in M$  :

$$\epsilon_x \cdot y = \begin{cases} \epsilon_x & \text{if } xy = x \\ 0 & \text{otherwise.} \end{cases}$$



# Simple modules

$M$  is  $\mathcal{J}$ -trivial and  $x \in M$ .

## Definition


Let  $S_x = \langle \epsilon_x \rangle$  vector space, right action of  $y \in M$  :

$$\epsilon_x \cdot y = \begin{cases} \epsilon_x & \text{if } xy = x \\ 0 & \text{otherwise.} \end{cases}$$


## Theorem (Denton, Hivert, Schilling, Thiéry)

*Any simple module is isomorphic to  $S_x$  for  $x \in M$  (thus one-dimensional).*


# Simple modules



# Simple modules




# Simple modules




# Simple modules : proof

- $\leq_{\mathcal{J}}$  partial order



## Simple modules : proof

- $\leq_{\mathcal{J}}$  partial order
- $(x_1, \dots, x_n)$  linear extension :  
 $x_i \leq_{\mathcal{J}} x_j \Rightarrow i \leq j$




# Simple modules : proof

- $\leq_{\mathcal{J}}$  partial order
- $(x_1, \dots, x_n)$  linear extension :  
 $x_i \leq_{\mathcal{J}} x_j \Rightarrow i \leq j$
- $F_i := \mathbb{K}\{x_j \mid j \leq i\}$ ,  $F_0 := \{0_{\mathbb{K}}\}$  :

$$F_0 \subset F_1 \subset \dots \subset F_{n-1} \subset F_n$$

composition series for  $F_n$  regular representation




# Simple modules : proof

- $\leq_{\mathcal{J}}$  partial order
- $(x_1, \dots, x_n)$  linear extension :  
 $x_i \leq_{\mathcal{J}} x_j \Rightarrow i \leq j$
- $F_i := \mathbb{K}\{x_j \mid j \leq i\}$ ,  $F_0 := \{0_{\mathbb{K}}\}$  :

$$F_0 \subset F_1 \subset \dots \subset F_{n-1} \subset F_n$$

composition series for  $F_n$  regular representation

- $F_i / F_{i-1} \cong S_{x_i}$





# Simple modules

Theorem (Norton, Carter & Denton, Hivert, Schilling, Thiéry)

$(S_e)_{e \in E(M)}$  complete set of pairwise non isomorphic simple modules  
Furthermore  $\{x - x^\omega \mid x \notin E(M)\}$  basis for  $\text{rad } \mathbb{K}M$ .

# Simple modules

Theorem (Norton, Carter & Denton, Hivert, Schilling, Thiéry)

$(S_e)_{e \in E(M)}$  complete set of pairwise non isomorphic simple modules  
 Furthermore  $\{x - x^\omega \mid x \notin E(M)\}$  basis for  $\text{rad } \mathbb{K}M$ .

Remark :


$$e \star f := (ef)^\omega$$

$$\mathbb{K}M / \text{rad } \mathbb{K}M \cong (\mathbb{K}E(M), \star)$$


Furthermore :

$$\begin{array}{ccc} \mathbb{K}M & \longrightarrow & \mathbb{K}M / \text{rad } \mathbb{K}M \\ x & \longmapsto & x^\omega \end{array}$$

# Simple modules



# Simple modules



Types :

- 000 : 1234
- 001 : 1243
- 010 : 1324
- 011 : 1432
- 100 : 2134
- 101 : 2143
- 110 : 3214
- 111 : 4321


$\epsilon_{3142}$

0

Type : 010

# Simple modules

$E(H_4^0)$  :



# Projective indecomposable modules

## Definition

Let  $M$  a  $\mathcal{J}$ -trivial monoid,  $x \in M$ . Then

$$\text{rAut}(x) := \{u \in M \mid xu = u\}.$$

# Projective indecomposable modules

## Definition

Let  $M$  a  $\mathcal{J}$ -trivial monoid,  $x \in M$ . Then

$$\text{rAut}(x) := \{u \in M \mid xu = u\}.$$

The smallest element of  $\text{rAut}(x)$  for the  $\mathcal{J}$ -order is called  $\text{rfix}(x)$  and

$$\text{rfix}(x) := \min\{e \in E(M) \mid xe = x\}.$$

# Projective indecomposable modules

## Definition

Let  $M$  a  $\mathcal{J}$ -trivial monoid,  $x \in M$ . Then

$$\text{rAut}(x) := \{u \in M \mid xu = u\}.$$

The smallest element of  $\text{rAut}(x)$  for the  $\mathcal{J}$ -order is called  $\text{rfix}(x)$  and

$$\text{rfix}(x) := \min\{e \in E(M) \mid xe = x\}.$$

Remark : a  $\mathcal{J}$ -trivial monoid always have a zero element.



# Projective indecomposable modules

## Definition

For  $x \in H_n^0$ , we define its right descent set by

$$D_R(x) = \{1 \leq i \leq n-1 \mid x\pi_i = x\}.$$

# Projective indecomposable modules

## Definition

For  $x \in H_n^0$ , we define its right descent set by

$$D_R(x) = \{1 \leq i \leq n-1 \mid x\pi_i = x\}.$$

## Example : positions

Let  $x = 413256$ .

# Projective indecomposable modules

## Definition

For  $x \in H_n^0$ , we define its right descent set by

$$D_R(x) = \{1 \leq i \leq n-1 \mid x\pi_i = x\}.$$

## Example : positions

Let  $x = 413256$ .  $D_R(x) = \{1, 3\}$ .

# Projective indecomposable modules

## Definition

For  $x \in H_n^0$ , we define its right descent set by

$$D_R(x) = \{1 \leq i \leq n-1 \mid x\pi_i = x\}.$$

## Example : positions

Let  $x = 413256$ .  $D_R(x) = \{1, 3\}$ .

Notation :

4			
1	3		
	2	5	6

# Projective indecomposable modules

## Definition

For  $x \in H_n^0$ , we define its right descent set by

$$D_R(x) = \{1 \leq i \leq n-1 \mid x\pi_i = x\}.$$

## Example : positions

Let  $x = 413256$ .  $D_R(x) = \{1, 3\}$ .

Notation :

4			
1	3		
	2	5	6

## Theorem (Denton, Hivert, Schilling, Thiéry)

$$\text{rfix}(x) = \pi_{D_R(x)}$$

# Projective indecomposable modules

## Definition

For  $x \in H_n^0$ , we define its right descent set by

$$D_R(x) = \{1 \leq i \leq n-1 \mid x\pi_i = x\}.$$

## Example : positions

Let  $x = 413256$ .  $D_R(x) = \{1, 3\}$ .

$\text{rfix}(x) = \pi_{\{1,3\}} = 214356$

Notation :

4					
1	3				
	2	5	6		

## Theorem (Denton, Hivert, Schilling, Thiéry)

$$\text{rfix}(x) = \pi_{D_R(x)}$$

# Projective indecomposable modules

Theorem (Norton, Carter & Denton, Hivert, Schilling, Thiéry)

$e \in E(M)$ . We set :

$$L(e) := Me$$

$$L_=(e) := \{x \in Me \mid \text{rfix}(x) = e\}$$

$$L_<(e) := \{x \in Me \mid \text{rfix}(x) <_{\mathcal{L}} e\}$$

# Projective indecomposable modules

Theorem (Norton, Carter & Denton, Hivert, Schilling, Thiéry)

$e \in E(M)$ . We set :

$$L(e) := Me$$

$$L_=(e) := \{x \in Me \mid \text{rfix}(x) = e\}$$

$$L_<(e) := \{x \in Me \mid \text{rfix}(x) <_{\mathcal{L}} e\}$$

Then, the projective module  $P_e$  associated to  $S_e$  is isomorphic to  $\mathbb{K}L(e) / \mathbb{K}L_<(e)$ .



# Projective indecomposable modules

Theorem (Norton, Carter & Denton, Hivert, Schilling, Thiéry)

$e \in E(M)$ . We set :

$$L(e) := Me$$

$$L_=(e) := \{x \in Me \mid \text{rfix}(x) = e\}$$


$$L_<(e) := \{x \in Me \mid \text{rfix}(x) <_{\mathcal{L}} e\}$$

Then, the projective module  $P_e$  associated to  $S_e$  is isomorphic to  $\mathbb{K}L(e)/\mathbb{K}L_<(e)$ .


Basis of  $P_e$  : image of  $L_=(e)$  in the quotient.

Action of  $m \in M$  on  $x \in L_=(e)$  :  $m \cdot x = \begin{cases} mx & \text{if } \text{rfix}(mx) = e \\ 0 & \text{otherwise.} \end{cases}$


# Projective indecomposable modules



# Projective indecomposable modules



# Projective indecomposable modules



# Effectivity

Simple modules, projective indecomposable modules, Cartan matrix, quiver.

# Effectivity

Simple modules, projective indecomposable modules, Cartan matrix, quiver.

- Algebra of finite dimension :

# Effectivity

Simple modules, projective indecomposable modules, Cartan matrix, quiver.

- Algebra of finite dimension :  $\dim A^6$

# Effectivity

Simple modules, projective indecomposable modules, Cartan matrix, quiver.

- Algebra of finite dimension :  $\dim A^6$
- Finite  $\mathcal{J}$ -trivial monoid :



# Effectivity

Simple modules, projective indecomposable modules, Cartan matrix, quiver.

- Algebra of finite dimension :  $\dim A^6$
- Finite  $\mathcal{J}$ -trivial monoid :  $|M|^3$

# Contents

- 1 Green Relations
- 2 The 0-Hecke monoid
- 3 Representation theory of  $\mathcal{J}$ -trivial monoid
- 4 The 0-rook monoid**
- 5 Renner monoids

# The rook monoid

**Rook matrix** of size  $n$  = set of non attacking rooks on an  $n \times n$  matrix.

# The rook monoid

**Rook matrix** of size  $n =$  set of non attacking rooks on an  $n \times n$  matrix.

$$\text{Rook Matrix} \quad \begin{pmatrix} 0 & 0 & 0 & 0 & \text{♚} \\ 0 & 0 & \text{♚} & 0 & 0 \\ 0 & 0 & 0 & \text{♚} & 0 \\ 0 & \text{♚} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & 0 & 0 & \text{♚} \\ 0 & 0 & 0 & 0 & 0 \\ 0 & \text{♚} & 0 & 0 & 0 \\ 0 & 0 & 0 & \text{♚} & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

# The rook monoid

**Rook matrix** of size  $n =$  set of non attacking rooks on an  $n \times n$  matrix.

$$\begin{array}{l}
 \text{Rook Matrix} \\
 \text{Rook Vector}
 \end{array}
 \begin{array}{cc}
 \begin{pmatrix} 0 & 0 & 0 & 0 & \text{♚} \\ 0 & 0 & \text{♚} & 0 & 0 \\ 0 & 0 & 0 & \text{♚} & 0 \\ 0 & \text{♚} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} & 
 \begin{pmatrix} 0 & 0 & 0 & 0 & \text{♚} \\ 0 & 0 & 0 & 0 & 0 \\ 0 & \text{♚} & 0 & 0 & 0 \\ 0 & 0 & 0 & \text{♚} & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \\
 \begin{pmatrix} 0 & 4 & 2 & 3 & 1 \end{pmatrix} & 
 \begin{pmatrix} 0 & 3 & 0 & 4 & 1 \end{pmatrix}
 \end{array}$$

# The rook monoid

**Rook matrix** of size  $n$  = set of non attacking rooks on an  $n \times n$  matrix.

$$\begin{array}{l}
 \text{Rook Matrix} \\
 \text{Rook Vector}
 \end{array}
 \begin{array}{c}
 \begin{pmatrix} 0 & 0 & 0 & 0 & \text{♚} \\ 0 & 0 & \text{♚} & 0 & 0 \\ 0 & 0 & 0 & \text{♚} & 0 \\ 0 & \text{♚} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \\
 \begin{pmatrix} 0 & 0 & 0 & 0 & \text{♚} \\ 0 & 0 & 0 & 0 & 0 \\ 0 & \text{♚} & 0 & 0 & 0 \\ 0 & 0 & 0 & \text{♚} & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}
 \end{array}
 \begin{array}{c}
 \begin{pmatrix} 0 & 0 & 0 & 0 & \text{♚} \\ 0 & 0 & 0 & 0 & 0 \\ 0 & \text{♚} & 0 & 0 & 0 \\ 0 & 0 & 0 & \text{♚} & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} \\
 \begin{pmatrix} 0 & 3 & 0 & 4 & 1 \end{pmatrix}
 \end{array}$$

The product of two rook matrices is a rook matrix.

**Rook Monoid**  $R_n$  = submonoid of the rook matrices

$$M_n \supset R_n \supset \mathfrak{S}_n$$

$\mathfrak{S}_n$

$$\mathfrak{S}_n \xleftarrow{q=1} \mathcal{H}_n(q)$$

# Iwahori



$$\mathfrak{S}_n \xleftarrow{q=1} \mathcal{H}_n(q) \xrightarrow{q=0} H_n^0$$

# Iwahori

$$\mathfrak{S}_n \xleftarrow{q=1} \mathcal{H}_n(q) \xrightarrow{q=0} H_n^0$$

$$\downarrow$$

$$R_n$$

# Iwahori

$$\mathfrak{S}_n \xleftarrow{q=1} \mathcal{H}_n(q) \xrightarrow{q=0} H_n^0$$



$$R_n \xleftarrow{q=1} \mathcal{I}_n(q)$$

Iwahori , Solomon.

$$\mathfrak{S}_n \xleftarrow{q=1} \mathcal{H}_n(q) \xrightarrow{q=0} H_n^0$$

$$\downarrow$$

$$\downarrow$$

$$\downarrow$$

$$R_n \xleftarrow{q=1} \mathcal{I}_n(q) \xrightarrow{q=0} ??$$

Iwahori , Solomon.

$$\begin{array}{ccccc}
 \mathfrak{S}_n & \xleftarrow{q=1} & \mathcal{H}_n(q) & \xrightarrow{q=0} & H_n^0 \\
 \downarrow & & \downarrow & & \downarrow \\
 R_n & \xleftarrow{q=1} & \mathcal{I}_n(q) & \xrightarrow{q=0} & R_n^0
 \end{array}$$

Iwahori , Solomon.

# Definition by right action on $R_n$

Operators  $\pi_0, \pi_1, \dots, \pi_{n-1}$  acting on rook vectors

# Definition by right action on $R_n$

Operators  $\pi_0, \pi_1, \dots, \pi_{n-1}$  acting on rook vectors

Bubble sort operators  $\pi_1, \dots, \pi_{n-1}$  :

$$(r_1 \dots r_n) \cdot \pi_i = \begin{cases} r_1 \dots r_{i-1} r_{i+1} r_i r_{i+2} \dots r_n & \text{if } r_i < r_{i+1}, \\ r_1 \dots r_n & \text{otherwise,} \end{cases}$$

Deletion operator  $\pi_0$  :

$$(r_1 \dots r_n) \cdot \pi_0 = 0 r_2 \dots r_n.$$

# Definition by right action on $R_n$

Operators  $\pi_0, \pi_1, \dots, \pi_{n-1}$  acting on rook vectors

Bubble sort operators  $\pi_1, \dots, \pi_{n-1}$  :

$$(r_1 \dots r_n) \cdot \pi_i = \begin{cases} r_1 \dots r_{i-1} r_{i+1} r_i r_{i+2} \dots r_n & \text{if } r_i < r_{i+1}, \\ r_1 \dots r_n & \text{otherwise,} \end{cases}$$

Deletion operator  $\pi_0$  :

$$(r_1 \dots r_n) \cdot \pi_0 = 0 r_2 \dots r_n.$$

$$45321 \cdot \pi_1 = 54321$$

$$40321 \cdot \pi_1 = 40321$$

$$00321 \cdot \pi_2 = 03021$$

$$3027006 \cdot \pi_0 = 0027006$$



# Definition by right action on $R_n$

Operators  $\pi_0, \pi_1, \dots, \pi_{n-1}$  acting on rook vectors


Bubble sort operators  $\pi_1, \dots, \pi_{n-1}$  :

$$(r_1 \dots r_n) \cdot \pi_i = \begin{cases} r_1 \dots r_{i-1} r_{i+1} r_i r_{i+2} \dots r_n & \text{if } r_i < r_{i+1}, \\ r_1 \dots r_n & \text{otherwise,} \end{cases}$$

Deletion operator  $\pi_0$  :

$$(r_1 \dots r_n) \cdot \pi_0 = 0 r_2 \dots r_n.$$

$$\begin{aligned} 45321 \cdot \pi_1 &= 54321 \\ 40321 \cdot \pi_1 &= 40321 \\ 00321 \cdot \pi_2 &= 03021 \\ 3027006 \cdot \pi_0 &= 0027006 \end{aligned}$$



# Definition by presentation

Generators :  $\pi_0, \dots, \pi_{n-1}$

Relations :

$$\pi_i^2 = \pi_i$$

$$0 \leq i \leq n-1,$$

$$\pi_i \pi_{i+1} \pi_i = \pi_{i+1} \pi_i \pi_{i+1}$$

$$1 \leq i \leq n-2,$$

$$\pi_i \pi_j = \pi_j \pi_i$$

$$|i-j| > 1.$$

$$\pi_1 \pi_0 \pi_1 \pi_0 = \pi_0 \pi_1 \pi_0 = \pi_0 \pi_1 \pi_0 \pi_1$$

# Definition by presentation

Generators :  $\pi_0, \dots, \pi_{n-1}$

Relations :

$$\pi_i^2 = \pi_i$$

$$\pi_i \pi_{i+1} \pi_i = \pi_{i+1} \pi_i \pi_{i+1}$$


$$\pi_i \pi_j = \pi_j \pi_i$$

$$\pi_1 \pi_0 \pi_1 \pi_0 = \pi_0 \pi_1 \pi_0 = \pi_0 \pi_1 \pi_0 \pi_1$$

$$0 \leq i \leq n-1,$$

$$1 \leq i \leq n-2,$$

$$|i-j| > 1.$$



## Theorem

*Both definitions (presentation and action on  $R_n$ ) are equivalent.*

## Theorem


*Both definitions (presentation and action on  $R_n$ ) are equivalent.*

Key Fact :

## Theorem

The map  $f : \left| \begin{array}{ll} R_n^0 & \longrightarrow R_n \\ r & \longmapsto \mathbf{1}_n \cdot r \end{array} \right. \text{ is a bijection.}$

# $\mathcal{J}$ -triviality : $R_n^0$ bisided Cayley graph



# Simple modules

## Theorem

$R_n^0$  is  $\mathcal{J}$ -trivial.

# Simple modules

## Theorem

$R_n^0$  is  $\mathcal{J}$ -trivial.

## Corollary (Application of Denton-Hivert-Schilling-Thiéry)

$R_n^0$  has  $2^n$  idempotents.

It has thus  $2^n$  simple modules of dimension 1.



# Descent set

## Definition

For  $\pi \in R_n^0$ , we define its right  $R$ -descent set by

$$D_R(\pi) = \{0 \leq i \leq n-1 \mid \pi\pi_i = \pi\}.$$

# Descent set

## Definition

For  $\pi \in R_n^0$ , we define its right  $R$ -descent set by

$$D_R(\pi) = \{0 \leq i \leq n-1 \mid \pi\pi_i = \pi\}.$$

## Example : positions

Let  $r = 0423007$ .  $0 < 4 \geq 2 < 3 \geq 0 \geq 0 < 7$ .

$$D_R(r) = \{0, 2, 4, 5\}$$

# Descent set

## Definition

For  $\pi \in R_n^0$ , we define its right  $R$ -descent set by

$$D_R(\pi) = \{0 \leq i \leq n-1 \mid \pi\pi_i = \pi\}.$$

## Example : positions

Let  $r = 0423007$ .  $0 < 4 \geq 2 < 3 \geq 0 \geq 0 < 7$ .

$D_R(r) = \{0, 2, 4, 5\}$  Notation :

0	4		
	2	3	
		0	
		0	7

# Descent set

## Definition

For  $\pi \in R_n^0$ , we define its right  $R$ -descent set by

$$D_R(\pi) = \{0 \leq i \leq n-1 \mid \pi\pi_i = \pi\}.$$

## Example : positions

Let  $r = 0423007$ .  $0 < 4 \geq 2 < 3 \geq 0 \geq 0 < 7$ .

$D_R(r) = \{0, 2, 4, 5\}$  Notation :

0	4		
	2	3	
		0	
		0	7

Warning : 

0
0


 and not 

0	0
---	---

.

# Descent class


List of the  $R$ -descent types for  $R_4^0$ :



# Projective modules

## Theorem (Application of Denton-Hivert-Schilling-Thiéry)

*The projective indecomposable  $R_n^0$ -modules are indexed by the  $R$ -descent type and isomorphic to the quotient of the associated  $R$ -descent class by the finer  $R$ -descent class.*



# Projectivity over $H_n^0$


## Theorem

*The indecomposable projective  $R_n^0$ -module splits as a  $H_n^0$ -module as the direct sum of all the indecomposable projective  $H_n^0$ -modules whose descent classes are explicit.*

Proof : explicit decomposition


$$\begin{array}{|c|c|} \hline 0 & \\ \hline \square & \square \\ \hline \end{array} = \begin{array}{|c|c|} \hline 0 & \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline 0 & \square \\ \hline \square & \square \\ \hline \end{array} = \begin{array}{|c|c|} \hline 0 & \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \\ \hline \square & \square \\ \hline \end{array} = \begin{array}{|c|c|} \hline \square & \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} = \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + 2 \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}.
 \end{array}$$

$$\begin{array}{|c|c|} \hline 0 \\ \hline \square \\ \hline \end{array} = \begin{array}{|c|c|} \hline 0 & \text{yellow} \\ \hline \text{yellow} & \text{yellow} \\ \hline \end{array} + \begin{array}{|c|c|} \hline 0 & \text{blue} \\ \hline \text{blue} & \text{blue} \\ \hline \end{array} = \begin{array}{|c|} \hline 0 \\ \hline \text{yellow} \\ \hline \text{yellow} \\ \hline \end{array} + \begin{array}{|c|} \hline 0 \\ \hline \text{blue} \\ \hline \text{blue} \\ \hline \end{array} = \begin{array}{|c|c|} \hline \text{yellow} & \text{yellow} \\ \hline \text{yellow} & \text{yellow} \\ \hline \end{array} + \begin{array}{|c|c|} \hline \text{yellow} & \text{yellow} \\ \hline \text{yellow} & \text{yellow} \\ \hline \end{array} + \begin{array}{|c|c|} \hline \text{blue} & \text{blue} \\ \hline \text{blue} & \text{blue} \\ \hline \end{array} + \begin{array}{|c|c|} \hline \text{blue} & \text{blue} \\ \hline \text{blue} & \text{blue} \\ \hline \end{array} + \begin{array}{|c|c|} \hline \text{blue} & \text{blue} \\ \hline \text{blue} & \text{blue} \\ \hline \end{array} + \begin{array}{|c|} \hline \text{blue} \\ \hline \text{blue} \\ \hline \text{blue} \\ \hline \end{array} = \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + 2 \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}.
 \end{array}$$






$$\begin{array}{|c|c|} \hline 0 \\ \hline \square \\ \hline \end{array} = \begin{array}{|c|c|} \hline 0 & \text{yellow} \\ \hline \text{yellow} & \text{yellow} \\ \hline \end{array} + \begin{array}{|c|c|} \hline 0 & \text{blue} \\ \hline \text{blue} & 0 \\ \hline \end{array} = \begin{array}{|c|} \hline 0 \\ \hline \text{yellow} \\ \hline \text{yellow} \\ \hline \end{array} + \begin{array}{|c|} \hline 0 \\ \hline \text{blue} \\ \hline \text{blue} \\ \hline \end{array} = \begin{array}{|c|} \hline \text{yellow} \\ \hline \text{yellow} \\ \hline \end{array} + \begin{array}{|c|c|} \hline \text{yellow} & \text{yellow} \\ \hline \text{yellow} & \text{yellow} \\ \hline \end{array} + \begin{array}{|c|c|} \hline \text{blue} & \text{blue} \\ \hline \text{blue} & \text{blue} \\ \hline \end{array} + \begin{array}{|c|c|} \hline \text{blue} & \text{blue} \\ \hline \text{blue} & \text{blue} \\ \hline \end{array} + \begin{array}{|c|} \hline \text{blue} \\ \hline \text{blue} \\ \hline \end{array} + \begin{array}{|c|} \hline \text{blue} \\ \hline \text{blue} \\ \hline \end{array} = \begin{array}{|c|c|} \hline & \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline & & \\ \hline \square & \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline & \\ \hline \square & \square \\ \hline \end{array} + 2 \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} + \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}.
 \end{array}$$



$$\begin{array}{|c|c|} \hline 0 \\ \hline \square \\ \hline \end{array} = \begin{array}{|c|c|} \hline 0 & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline 0 & \square \\ \hline \square & 0 \\ \hline \end{array} = \begin{array}{|c|} \hline 0 \\ \hline \square \\ \hline \square \\ \hline \end{array} + \begin{array}{|c|} \hline 0 \\ \hline \square \\ \hline \square \\ \hline \end{array} = \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} = \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array} + 2 \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array} + \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}.
 \end{array}$$



# Tower of monoids

$$R_1^0 \subset R_2^0 \subset \dots \subset R_n^0 \subset \dots$$

# Tower of monoids

$$R_1^0 \subset R_2^0 \subset \cdots \subset R_n^0 \subset \cdots$$

Results :

- Induction of simple and projective modules

# Tower of monoids

$$R_1^0 \subset R_2^0 \subset \cdots \subset R_n^0 \subset \cdots$$

Results :

- Induction of simple and projective modules
- Restriction of simple modules

# Tower of monoids

$$R_1^0 \subset R_2^0 \subset \cdots \subset R_n^0 \subset \cdots$$

Results :

- Induction of simple and projective modules
- Restriction of simple modules
- Restriction of projective modules : does not work

# Contents

- 1 Green Relations
- 2 The 0-Hecke monoid
- 3 Representation theory of  $\mathcal{J}$ -trivial monoid
- 4 The 0-rook monoid
- 5 Renner monoids**

# Algebraic monoid

## Definition

An *algebraic monoid* is a submonoid of  $M_n$  closed for the Zariski topology.



# Algebraic monoid

## Definition

An *algebraic monoid* is a submonoid of  $M_n$  closed for the Zariski topology.

If  $G$  algebraic subgroup of  $GL_n$  then  $M = \overline{G}$  algebraic monoid with  $G$  as unit group.

If  $M$  algebraic monoid,  $G(M)$  unit group is algebraic group.

# Algebraic monoid

## Definition

An *algebraic monoid* is a submonoid of  $M_n$  closed for the Zariski topology.

If  $G$  algebraic subgroup of  $GL_n$  then  $M = \overline{G}$  algebraic monoid with  $G$  as unit group.

If  $M$  algebraic monoid,  $G(M)$  unit group is algebraic group.

## Definition

$M$  is regular if  $M = E(M)G(M) = G(M)E(M)$ .

# Algebraic monoid

## Definition

An *algebraic monoid* is a submonoid of  $M_n$  closed for the Zariski topology.

If  $G$  algebraic subgroup of  $GL_n$  then  $M = \overline{G}$  algebraic monoid with  $G$  as unit group.

If  $M$  algebraic monoid,  $G(M)$  unit group is algebraic group.

## Definition

$M$  is regular if  $M = E(M)G(M) = G(M)E(M)$ .

## Theorem

$M$  irreducible algebraic monoid with a zero element.

$M$  is regular iff  $G(M)$  is reductive.

# Renner monoids and Weyl groups

## Definition

$M$  regular irreducible algebraic monoid with a zero element.

$T$  maximal Torus inside a Borel subgroup of  $G(M)$ , its normalizer  $N_{G(M)}(T)$ .

The Renner monoid  $R(M)$  is  $\overline{N_{G(M)}(T)} / T$ .

# Renner monoids and Weyl groups

## Definition

$M$  regular irreducible algebraic monoid with a zero element.

$T$  maximal Torus inside a Borel subgroup of  $G(M)$ , its normalizer  $N_{G(M)}(T)$ .

The Renner monoid  $R(M)$  is  $\overline{N_{G(M)}(T)} / T$ .

$W := N_{G(M)}(T) / T$  Weyl Group

$W$  is the unit group of  $R(M)$

$R(M) = E(\overline{T}) \cdot W = W \cdot E(\overline{T})$

# Renner monoids and Weyl groups

## Definition

$M$  regular irreducible algebraic monoid with a zero element.

$T$  maximal Torus inside a Borel subgroup of  $G(M)$ , its normalizer  $N_{G(M)}(T)$ .

The Renner monoid  $R(M)$  is  $\overline{N_{G(M)}(T)} / T$ .

$W := N_{G(M)}(T) / T$  Weyl Group

$W$  is the unit group of  $R(M)$

$$R(M) = E(\overline{T}) \cdot W = W \cdot E(\overline{T})$$

## Example

$M = M_n$  then  $R(M) = R_n$

# Results

- Definition of 0-Renner monoids in type  $B, D$ 
  - Presentation
  - Function on some rook vectors

# Results

- Definition of 0-Renner monoids in type  $B, D$ 
  - Presentation
  - Function on some rook vectors

Example : type  $B_5$

$1_{10}$  :

$\bar{5}\bar{4}\bar{3}\bar{2}\bar{1}12345$



# Results

- Definition of 0-Renner monoids in type  $B, D$ 
  - Presentation
  - Function on some rook vectors

Example : type  $B_5$

$1_{10} \cdot \pi_1 :$

$\bar{5}\bar{4}\bar{3}\bar{2}\bar{1}12345$

# Results

- Definition of 0-Renner monoids in type  $B, D$ 
  - Presentation
  - Function on some rook vectors

Example : type  $B_5$

$1_{10} \cdot \pi_1 :$

$\bar{5}\bar{4}\bar{3}\bar{2}\bar{1}12345$

# Results

- Definition of 0-Renner monoids in type  $B, D$ 
  - Presentation
  - Function on some rook vectors

Example : type  $B_5$

$1_{10} \cdot \pi_1 \pi_3 :$

$\bar{5}\bar{4}\bar{3}\bar{2}\bar{1}\bar{1}2345$

# Results

- Definition of 0-Renner monoids in type  $B, D$ 
  - Presentation
  - Function on some rook vectors

Example : type  $B_5$

$1_{10} \cdot \pi_1 \pi_3 :$

$\bar{5} \bar{3} \bar{4} \bar{2} \bar{1} \bar{1} \bar{2} 4 3 5$

# Results

- Definition of 0-Renner monoids in type  $B, D$ 
  - Presentation
  - Function on some rook vectors

Example : type  $B_5$

$1_{10} \cdot \pi_1 \pi_3 \pi_0 :$

$\bar{5}\bar{3}\bar{4}\bar{2}\bar{1}\bar{1}2435$

# Results

- Definition of 0-Renner monoids in type  $B, D$ 
  - Presentation
  - Function on some rook vectors

Example : type  $B_5$

$1_{10} \cdot \pi_1 \pi_3 \pi_0 :$

00000 $\bar{1}$ 2435

# Results


- Definition of 0-Renner monoids in type  $B$ ,  $D$ 
  - Presentation
  - Function on some rook vectors
- Equivalence of the definitions

# Results


- Definition of 0-Renner monoids in type  $B, D$ 
  - Presentation
  - Function on some rook vectors
- Equivalence of the definitions
- On going :  $\mathcal{J}$ -triviality



# Results



# Results



# Results

- Definition of 0-Renner monoids in type  $B, D$ 
  - Presentation
  - Function on some rook vectors
- Equivalence of the definitions
- On going :  $\mathcal{J}$ -triviality
- On going : representation theory

**THANK YOU FOR YOUR  
OUTSTANDING  
ATTENTION!!**

# Descent classes are not intervals

