



Garside structures for $B(e, e, n)$

Georges Neaime

Laboratoire de Mathématiques Nicolas Oresme



Paris, April 27, 2017

Complex reflection groups

An element $s \in GL_n(\mathbb{C})$ is called a **reflection** if

- $\text{Ker}(s - 1)$ is a hyperplane,
- $s^2 = 1$.

Complex reflection groups

An element $s \in GL_n(\mathbb{C})$ is called a **reflection** if

- $\text{Ker}(s - 1)$ is a hyperplane,
- $s^2 = 1$.

Relaxing the second condition to s of finite order defines the notion of **pseudo-reflection**.

Complex reflection groups

An element $s \in GL_n(\mathbb{C})$ is called a **reflection** if

- $\text{Ker}(s - 1)$ is a hyperplane,
- $s^2 = 1$.

Relaxing the second condition to s of finite order defines the notion of **pseudo-reflection**.

Let W be a finite subgroup of $GL_n(\mathbb{C})$.

W is a **reflection group** if $W = \langle \mathcal{R} \rangle$ where \mathcal{R} is the set of reflections of W .

Complex reflection groups

An element $s \in GL_n(\mathbb{C})$ is called a **reflection** if

- $\text{Ker}(s - 1)$ is a hyperplane,
- $s^2 = 1$.

Relaxing the second condition to s of finite order defines the notion of **pseudo-reflection**.

Let W be a finite subgroup of $GL_n(\mathbb{C})$.

W is a **reflection group** if $W = \langle \mathcal{R} \rangle$ where \mathcal{R} is the set of reflections of W .

Fact: Every reflection group is a direct product of irreducible ones.

Irreducible reflection groups

The Shephard and Todd classification:

Irreducible reflection groups

The Shephard and Todd classification:

The first family:

For $e, n \geq 1$, $G(e, e, n)$ is the group of $n \times n$ matrices consisting of:

- monomial matrices (each row and column has a unique nonzero entry),
- with all nonzero entries lying in μ_e , the e -th roots of unity, and
- for which the product of the nonzero entries is 1.

Irreducible reflection groups

The second family:

For $e, n \geq 1$, $G(2e, e, n)$ is the group of $n \times n$ matrices consisting of:

- monomial matrices,
- with all nonzero entries lying in μ_{2e} , and
- for which the product of the nonzero entries is 1 or -1 .

Irreducible reflection groups

The second family:

For $e, n \geq 1$, $G(2e, e, n)$ is the group of $n \times n$ matrices consisting of:

- monomial matrices,
- with all nonzero entries lying in μ_{2e} , and
- for which the product of the nonzero entries is 1 or -1 .

Plus: 15 exceptions!

Complex braid groups

Let $W < GL_n(\mathbb{C})$ a complex reflection group, and \mathcal{R} the set of reflections of W .

Complex braid groups

Let $W < GL_n(\mathbb{C})$ a complex reflection group, and \mathcal{R} the set of reflections of W .

There is a corresponding hyperplane arrangement and hyperplane complement:

$$\mathcal{A} = \{ \text{Ker}(s - 1) \mid s \in \mathcal{R} \}, X = \mathbb{C}^n \setminus \cup \mathcal{A}$$

Complex braid groups

Let $W < GL_n(\mathbb{C})$ a complex reflection group, and \mathcal{R} the set of reflections of W .

There is a corresponding hyperplane arrangement and hyperplane complement:

$$\mathcal{A} = \{ \text{Ker}(s - 1) \mid s \in \mathcal{R} \}, X = \mathbb{C}^n \setminus \bigcup \mathcal{A}$$

- $P = \pi_1(X)$ pure (complex) braid group,
- $B = \pi_1(X/W)$ **braid group**, with

$$1 \longrightarrow P \longrightarrow B \longrightarrow W \longrightarrow 1$$

Artin-Tits groups

A finite **Coxeter** group W is defined by the following presentation:

$$W = \langle S \mid s^2 = 1, \underbrace{sts \cdots}_{m_{st}} = \underbrace{tst \cdots}_{m_{st}} \text{ for } s \neq t \in S \rangle$$

Artin-Tits groups

A finite **Coxeter** group W is defined by the following presentation:

$$W = \langle S \mid s^2 = 1, \underbrace{sts \cdots}_{m_{st}} = \underbrace{tst \cdots}_{m_{st}} \text{ for } s \neq t \in S \rangle$$

The **Artin-Tits** group $B(W)$ attached to W is the group of fractions of the monoid $B^+(W)$:

$$B^+(W) = \langle \tilde{S} \mid \underbrace{\tilde{s}\tilde{t}\tilde{s} \cdots}_{m_{st}} = \underbrace{\tilde{t}\tilde{s}\tilde{t} \cdots}_{m_{st}} \text{ for } \tilde{s} \neq \tilde{t} \in \tilde{S} \rangle^+$$

$$B(W) = \langle \tilde{S} \mid \underbrace{\tilde{s}\tilde{t}\tilde{s} \cdots}_{m_{st}} = \underbrace{\tilde{t}\tilde{s}\tilde{t} \cdots}_{m_{st}} \text{ for } \tilde{s} \neq \tilde{t} \in \tilde{S} \rangle$$

Link with the first definitions

Let $W < GL_n(\mathbb{R}) < GL_n(\mathbb{C})$.

Link with the first definitions

Let $W < GL_n(\mathbb{R}) < GL_n(\mathbb{C})$.

W a **real** reflection group

(Coxeter) \updownarrow

W is a **Coxeter** group:
 Generators + quadratic
 and braid relations

\rightsquigarrow

B the **Artin-Tits** group:
 Generators + braid relations

\updownarrow (Brieskorn)

$$B = \pi_1(X/W)$$

General goal

General goal: Extend what is known for Artin-Tits groups to complex braid groups.

General goal

General goal: Extend what is known for Artin-Tits groups to complex braid groups.

For example:

- Artin-Tits groups are **linear** (Krammer, Bigelow, Digne, Cohen-Wales, Paris)
- Artin-Tits groups admit **Garside** structures (Garside, Deligne)

General goal

General goal: Extend what is known for Artin-Tits groups to complex braid groups.

For example:

- Artin-Tits groups are **linear** (Krammer, Bigelow, Digne, Cohen-Wales, Paris)
- Artin-Tits groups admit **Garside** structures (Garside, Deligne)

Why considering $B(e, e, n)$, the complex braid group attached to $G(e, e, n)$?

Garside structure

Let us recall the notion of a Garside structure.

Garside structure

Let us recall the notion of a Garside structure.

Let M be a monoid and $f, g \in M$.

We say that f **left-divides** g , written $f \preceq g$, if $fg' = g$ holds for some $g' \in M$.

Similarly, we can define the notion of right division.

Garside structure

Definition

We say that M is a **Garside monoid** if

- M is cancellative,
- there exists $\lambda : M \rightarrow \mathbb{N}$ s.t. $\lambda(fg) \geq \lambda(f) + \lambda(g)$ and $g \neq 1 \implies \lambda(g) \neq 0$,
- any two elements of M have a gcd and an lcm,
- there exists an element $\Delta \in M$ such that its left and right divisors coincide and generate M , and the set of all divisors of Δ is finite.

Garside structure

Definition

We say that M is a **Garside monoid** if

- M is cancellative,
- there exists $\lambda : M \rightarrow \mathbb{N}$ s.t. $\lambda(fg) \geq \lambda(f) + \lambda(g)$ and $g \neq 1 \implies \lambda(g) \neq 0$,
- any two elements of M have a gcd and an lcm,
- there exists an element $\Delta \in M$ such that its left and right divisors coincide and generate M , and the set of all divisors of Δ is finite.

The element Δ is called a **Garside element** of M and the divisors of Δ are called the **simples** of the Garside structure.

Garside structure

Definition

We say that M is a **Garside monoid** if

- M is cancellative,
- there exists $\lambda : M \rightarrow \mathbb{N}$ s.t. $\lambda(fg) \geq \lambda(f) + \lambda(g)$ and $g \neq 1 \implies \lambda(g) \neq 0$,
- any two elements of M have a gcd and an lcm,
- there exists an element $\Delta \in M$ such that its left and right divisors coincide and generate M , and the set of all divisors of Δ is finite.

The element Δ is called a **Garside element** of M and the divisors of Δ are called the **simples** of the Garside structure.

A **Garside group** is the group of fractions of a Garside monoid.

Interval structure

Let G be a finite group generated by a set S .

Interval structure

Let G be a finite group generated by a set S .

We define a partial order relation on G by:

$$x \preceq y \iff \ell(x) + \ell(x^{-1}y) = \ell(y).$$

Interval structure

Let G be a finite group generated by a set S .

We define a partial order relation on G by:

$$x \preceq y \iff \ell(x) + \ell(x^{-1}y) = \ell(y).$$

For $g \in G$, we define a monoid $M([1, g])$ by the presentation with

- Generators: A set \underline{P} in bijection with the interval

$$[1, g] := \{x \in G \mid 1 \preceq x \preceq g\}$$

- Relations: $\underline{x} \underline{y} = \underline{z}$ if $xy = z$ and $x \preceq z \preceq g$, that is $\ell(x) + \ell(y) = \ell(z)$.

Interval structure

Similarly, we can introduce the partial order relation

$x \preceq_r y \iff \ell(yx^{-1}) + \ell(x) = \ell(y)$, and the interval $[1, g]_r$ then define the monoid $M([1, g]_r)$.

Interval structure

Similarly, we can introduce the partial order relation $x \preceq_r y \iff \ell(yx^{-1}) + \ell(x) = \ell(y)$, and the interval $[1, g]_r$ then define the monoid $M([1, g]_r)$.

Theorem (Michel)

If $[1, g] = [1, g]_r$ and both posets $([1, g], \preceq)$ and $([1, g]_r, \preceq_r)$ are lattices, then $M([1, g])$ is a **Garside** monoid with Garside element \underline{g} and with simples \underline{P} .

Interval structure

Seminal example:

Interval structure

Seminal example:

Let W be a finite coxeter group

$$W = \langle S \mid s^2 = 1, \underbrace{sts \cdots}_{m_{st}} = \underbrace{tst \cdots}_{m_{st}} \text{ for } s \neq t \in S \rangle$$

Interval structure

Seminal example:

Let W be a finite coxeter group

$$W = \langle S \mid s^2 = 1, \underbrace{sts \cdots}_{m_{st}} = \underbrace{tst \cdots}_{m_{st}} \text{ for } s \neq t \in S \rangle$$

Take $G = W$ and $g = w_0$ the longest element.

Interval structure

Seminal example:

Let W be a finite coxeter group

$$W = \langle S \mid s^2 = 1, \underbrace{sts \cdots}_{m_{st}} = \underbrace{tst \cdots}_{m_{st}} \text{ for } s \neq t \in S \rangle$$

Take $G = W$ and $g = w_0$ the longest element.

We have $[1, w_0] = W$.

Interval structure

Seminal example:

Let W be a finite coxeter group

$$W = \langle S \mid s^2 = 1, \underbrace{sts \cdots}_{m_{st}} = \underbrace{tst \cdots}_{m_{st}} \text{ for } s \neq t \in S \rangle$$

Take $G = W$ and $g = w_0$ the longest element.

We have $[1, w_0] = W$.

The interval monoid $M([1, w_0])$ is the Artin-Tits monoid $B^+(W)$.

Interval structure

Seminal example:

Let W be a finite coxeter group

$$W = \langle S \mid s^2 = 1, \underbrace{sts \cdots}_{m_{st}} = \underbrace{tst \cdots}_{m_{st}} \text{ for } s \neq t \in S \rangle$$

Take $G = W$ and $g = w_0$ the longest element.

We have $[1, w_0] = W$.

The interval monoid $M([1, w_0])$ is the Artin-Tits monoid $B^+(W)$.

Hence $B^+(W)$ is generated by a copy \underline{W} of W with $\underline{x} \underline{y} = \underline{z}$ if $xy = z$ and $\ell(x) + \ell(y) = \ell(z)$; $x, y, z \in W$.

Interval structure

Seminal example:

Let W be a finite coxeter group

$$W = \langle S \mid s^2 = 1, \underbrace{sts \cdots}_{m_{st}} = \underbrace{tst \cdots}_{m_{st}} \text{ for } s \neq t \in S \rangle$$

Take $G = W$ and $g = w_0$ the longest element.

We have $[1, w_0] = W$.

The interval monoid $M([1, w_0])$ is the Artin-Tits monoid $B^+(W)$.

Hence $B^+(W)$ is generated by a copy \underline{W} of W with $\underline{x} \underline{y} = \underline{z}$ if $xy = z$ and $\ell(x) + \ell(y) = \ell(z)$; $x, y, z \in W$.

Theorem

$B^+(W)$ is a **Garside monoid** with Garside element $\underline{w_0}$ and with simples \underline{W} .

Definition of $G(e, e, n)$

Recall that

Definition of $G(e, e, n)$

Recall that

for $e, n \geq 1$, $G(e, e, n)$ is the group of $n \times n$ matrices consisting of:


- monomial matrices,
- with all nonzero entries lying in μ_e , the e -th roots of unity, and
- for which the product of the nonzero entries is 1.

Presentation for $G(e, e, n)$

Define a group by a **presentation** with generators and relations that can be described by the following diagram:

Presentation for $G(e, e, n)$

Define a group by a **presentation** with generators and relations that can be described by the following diagram:



Presentation for $G(e, e, n)$

It is shown that this group is **isomorphic** to the complex reflection group $G(e, e, n)$ (Broué-Malle-Rouquier),

$$\text{with } \mathbf{t}_i \longmapsto t_i := \begin{pmatrix} 0 & \zeta_e^{-i} & 0 \\ \zeta_e^i & 0 & 0 \\ 0 & 0 & I_{n-2} \end{pmatrix} \text{ with } i = 0, 1 \text{ and}$$

$$\mathbf{s}_j \longmapsto s_j := \begin{pmatrix} I_{j-2} & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & I_{n-j} \end{pmatrix} \text{ with } 3 \leq j \leq n.$$

Presentation for $G(e, e, n)$

It is shown that this group is **isomorphic** to the complex reflection group $G(e, e, n)$ (Broué-Malle-Rouquier),

$$\text{with } \mathbf{t}_i \longmapsto t_i := \begin{pmatrix} 0 & \zeta_e^{-i} & 0 \\ \zeta_e^i & 0 & 0 \\ 0 & 0 & I_{n-2} \end{pmatrix} \text{ with } i = 0, 1 \text{ and}$$

$$\mathbf{s}_j \longmapsto s_j := \begin{pmatrix} I_{j-2} & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & I_{n-j} \end{pmatrix} \text{ with } 3 \leq j \leq n.$$


Coxeter cases $n = 2$ (dihedral groups), $e = 1$ (symmetric groups),
 $e = 2$ (type D_n).

Presentation of B-M-R for $B(e, e, n)$


Removing the quadratic relations from the presentation of Broué-Malle-Rouquier gives a **presentation** of the complex braid group $B(e, e, n)$ attached to $G(e, e, n)$ with diagram:

Presentation of B-M-R for $B(e, e, n)$

Removing the quadratic relations from the presentation of Broué-Malle-Rouquier gives a **presentation** of the complex braid group $B(e, e, n)$ attached to $G(e, e, n)$ with diagram:




Presentation of B-M-R for $B(e, e, n)$



This is called the **presentation of Broué-Malle-Rouquier** of $B(e, e, n)$.


Presentation of B-M-R for $B(e, e, n)$



This is called the **presentation of Broué-Malle-Rouquier** of $B(e, e, n)$.

If $e \geq 3$ and $n \geq 3$, the monoid defined by this presentation **fails to embed** in $B(e, e, n)$ (Corran)

Presentation of B-M-R for $B(e, e, n)$



This is called the **presentation of Broué-Malle-Rouquier** of $B(e, e, n)$.

If $e \geq 3$ and $n \geq 3$, the monoid defined by this presentation **fails to embed** in $B(e, e, n)$ (Corran)


Thus, this presentation does not give rise to a Garside structure for $B(e, e, n)$!

Presentation of C-P for $B(e, e, n)$

Let $\tilde{t}_i := \tilde{t}_{i-1}\tilde{t}_{i-2}\tilde{t}_{i-1}^{-1}$ for $2 \leq i \leq e-1$. Consider the following diagram **presentation**:


Presentation of C-P for $B(e, e, n)$

Let $\tilde{t}_i := \tilde{t}_{i-1}\tilde{t}_{i-2}\tilde{t}_{i-1}^{-1}$ for $2 \leq i \leq e-1$. Consider the following diagram **presentation**:



Presentation of C-P for $B(e, e, n)$


Let $\tilde{t}_i := \tilde{t}_{i-1}\tilde{t}_{i-2}\tilde{t}_{i-1}^{-1}$ for $2 \leq i \leq e-1$. Consider the following diagram **presentation**:



It is called the **presentation of Corran-Picantin** of $B(e, e, n)$.

Presentation of C-P for $B(e, e, n)$

Let $\tilde{t}_i := \tilde{t}_{i-1}\tilde{t}_{i-2}\tilde{t}_{i-1}^{-1}$ for $2 \leq i \leq e-1$. Consider the following diagram **presentation**:




It is called the **presentation of Corran-Picantin** of $B(e, e, n)$.

This presentation gives rise to a **Garside** structure for $B(e, e, n)$!

Presentation for $G(e, e, n)$

Adding $x^2 = 1$ for all generators x of the presentation of Corran-Picantin gives a **presentation** of a group isomorphic to $G(e, e, n)$



Presentation for $G(e, e, n)$

Adding $x^2 = 1$ for all generators x of the presentation of Corran-Picantin gives a **presentation** of a group isomorphic to $G(e, e, n)$,

Presentation for $G(e, e, n)$

Adding $x^2 = 1$ for all generators x of the presentation of Corran-Picantin gives a **presentation** of a group isomorphic to $G(e, e, n)$,

$$\text{with } \mathbf{t}_i \mapsto t_i := \begin{pmatrix} 0 & \zeta_e^{-i} & 0 \\ \zeta_e^i & 0 & 0 \\ 0 & 0 & I_{n-2} \end{pmatrix} \text{ with } 0 \leq i \leq e-1 \text{ and}$$

$$\mathbf{s}_j \mapsto s_j := \begin{pmatrix} I_{j-2} & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & I_{n-j} \end{pmatrix} \text{ with } 3 \leq j \leq n.$$

Simples and Garside element

Consider the presentation of Corran and Picantin of $B(e, e, n)$ which gives rise to a Garside structure.

Simples and Garside element

Consider the presentation of Corran and Picantin of $B(e, e, n)$ which gives rise to a Garside structure.

The **Garside element** is $\Delta = \underbrace{\tilde{t}_1 \tilde{t}_0}_{\Delta_2} \underbrace{\tilde{s}_3 \tilde{t}_1 \tilde{t}_0 \tilde{s}_3}_{\Delta_3} \cdots \underbrace{\tilde{s}_n \cdots \tilde{s}_3 \tilde{t}_1 \tilde{t}_0 \tilde{s}_3 \cdots \tilde{s}_n}_{\Delta_n},$

and

Simples and Garside element

Consider the presentation of Corran and Picantin of $B(e, e, n)$ which gives rise to a Garside structure.

The **Garside element** is $\Delta = \underbrace{\tilde{t}_1 \tilde{t}_0}_{\Delta_2} \underbrace{\tilde{s}_3 \tilde{t}_1 \tilde{t}_0 \tilde{s}_3}_{\Delta_3} \cdots \underbrace{\tilde{s}_n \cdots \tilde{s}_3 \tilde{t}_1 \tilde{t}_0 \tilde{s}_3 \cdots \tilde{s}_n}_{\Delta_n},$

and

the **simples** are precisely the elements of the form $\delta_2 \delta_3 \cdots \delta_n$ where δ_i is a divisor of Δ_i for $2 \leq i \leq n$.

Second homology group

Theorem (Callegaro, Marin)

Let $B := B(e, e, n)$ with $e \geq 2$ and $n \geq 2$.

- If $n = 2$, then $H_2(B, \mathbb{Z}) = 0$ if e is odd and $H_2(B, \mathbb{Z}) = \mathbb{Z}$ if e is even (Salvetti).
- If $n = 3$, then $H_2(B, \mathbb{Z}) \simeq \mathbb{Z}/e\mathbb{Z}$.
- If $n = 4$ and e is odd, then $H_2(B, \mathbb{Z}) \simeq \mathbb{Z}/e\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$.
- If $n = 4$ and e is even, then $H_2(B, \mathbb{Z}) \simeq \mathbb{Z}/e\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^2$.
- If $n \geq 5$, then $H_2(B, \mathbb{Z}) \simeq \mathbb{Z}/e\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$.

Parabolic submonoids

Let δ be a balanced element of $Div(\Delta)$. Denote by B_δ the subgroup of $B(e, e, n)$ generated by the elements of $\{\tilde{t}_0, \tilde{t}_1, \dots, \tilde{t}_{e-1}, \tilde{s}_3, \dots, \tilde{s}_n\}$ that are in $Div(\delta)$.

Parabolic submonoids

Let δ be a balanced element of $Div(\Delta)$. Denote by B_δ the subgroup of $B(e, e, n)$ generated by the elements of $\{\tilde{t}_0, \tilde{t}_1, \dots, \tilde{t}_{e-1}, \tilde{s}_3, \dots, \tilde{s}_n\}$ that are in $Div(\delta)$.

We say that B_δ is a **standard parabolic subgroup** of $B(e, e, n)$ if $Div(\delta) = Div(\Delta) \cap B_\delta$ (definition of Godelle for standard parabolic subgroups in a Garside structure).

Parabolic submonoids


Let δ be a balanced element of $Div(\Delta)$. Denote by B_δ the subgroup of $B(e, e, n)$ generated by the elements of $\{\tilde{t}_0, \tilde{t}_1, \dots, \tilde{t}_{e-1}, \tilde{s}_3, \dots, \tilde{s}_n\}$ that are in $Div(\delta)$.

We say that B_δ is a **standard parabolic subgroup** of $B(e, e, n)$ if $Div(\delta) = Div(\Delta) \cap B_\delta$ (definition of Godelle for standard parabolic subgroups in a Garside structure).

The balanced elements of $Div(\Delta)$ are the lcm of the subsets of $\{\tilde{t}_0, \tilde{t}_1, \dots, \tilde{t}_{e-1}, \tilde{s}_3, \dots, \tilde{s}_n\}$. Denote δ one of these balanced elements. We have B_δ a standard parabolic subgroup of $B(e, e, n)$ in the sense of Godelle.


Parabolic submonoids

For example, consider the presentation of Corran-Picantin for $B(e, e, 3)$



Parabolic submonoids


For example, consider the presentation of Corran-Picantin for $B(e, e, 3)$





We have **2 types** of standard parabolic subgroups in the sense of Godelle:


Parabolic submonoids

For example, consider the presentation of Corran-Picantin for $B(e, e, 3)$



We have **2 types** of standard parabolic subgroups in the sense of Godelle:





Thank you!